

addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

Amendments

In the Title:

Please substitute the following Title of the Invention for the pending Title of the Invention:


Aryl Substituted Pyrimidines

In the Claims:


Please cancel claims 13, 16, 27, 30, 40, 49, and 52-58 without prejudice or disclaimer.

Please substitute the following claim 1 for the pending claim 1:

1. (Once amended) A compound having the Formula I:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:



Y is or R₇,

provided that when Y is R₇, R₁ is aminocarbonyl;

BT
Sub
C1

A₁ is N and A₂ and A₃ are CR₂, or A₃ is N and A₁ and A₂ are CR₂;

R₁ is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, C(O)R₈, SO₂R₈, OC(O)NH₂, 2-imidazolyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

Sub C1
each R₂ is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, amino, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

B1
R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₇ is an optionally substituted alkyl;

R₈ is selected from the group consisting of alkyl, alkenyl, alkynyl, OR₉, amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R₈ is not OR₉ when R₁ is SO₂R₈; wherein

R₉ is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and


X is one of O, S, NH, or CH₂ when Y is other than R₇; or

X is one of O, S, NH, CH₂ or absent when Y is R₇;

with the proviso that R₂ is not methoxy if R₅ is trifluoromethyl, R₆ is H, X is O and R₁ is SO₂CH₂Ph.

Please substitute the following claim 2 for the pending claim 2:

2. (Once Amended) A compound having the Formula II:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:

A₁ is N and A₂ and A₃ are CR₂, or A₃ is N and A₁ and A₂ are CR₂;

R₁ is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, C(O)R₈, SO₂R₈, OC(O)NH₂, 2-imidazoliny, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R₂ is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, amino, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol; and

R₈ is selected from the group consisting of alkyl, alkenyl, alkynyl, OR₉, amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R₈ is not OR₉ when R₁ is SO₂R₈; wherein

R₉ is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and

X is one of O, S, NH, or CH₂;

Sub
C2

B2

Sub
C2 B2 with the proviso that R₂ is not methoxy if R₅ is trifluoromethyl, R₆ is H, X is O
and R₁ is SO₂CH₂Ph.

Please substitute the following claim 3 for the pending claim 3:

D4
B3 [3. (Once Amended) The compound of claim 2, wherein A₃ is N and A₁ and A₂
are CR₂.

Please substitute the following claim 18 for the pending claim 18:

D4
B4 [18. (Once Amended) The compound of claim 17, wherein A₃ is N and A₁ and A₂
are CR₂.

Please substitute the following claim 26 for the pending claim 26:

D4
B5 [26. (Once Amended) The compound of claim 17, wherein
X is O;
A₁ is N and A₂ and A₃ are CR₂; or A₃ is N and A₁ and A₂ are CR₂; wherein
R₂ is selected from the group consisting of hydrogen, alkyl, alkoxy, aminoalkyl,
and aminocarbonyl;
R₃ and R₄ are both hydrogen;
R₅ and R₆ are independently selected from the group consisting of hydrogen,
alkyl, halogen, haloalkyl, and nitro; and
R₈ is amino.

Please substitute the following claim 32 for the pending claim 32:

D4
B6 [32. (Once Amended) The compound of claim 31, wherein R₂ is selected from
the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aminoalkyl, amino,
hydroxyalkyl, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl,
aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and
aralkylcarbonylamino.

Please substitute the following claim 39 for the pending claim 39:

39. (Once Amended) A compound of claim 2, wherein said compound is:

- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(4-nitrophenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(4-methoxyphenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(4-trifluoromethylphenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(3-chloro-2-cyanophenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(2,4-difluorophenoxy)phenyl]pyrimidine-2-carboxamide;
- 4-[4-(2-chloro-4-fluorophenoxy)phenyl]pyrimidine-2-carboxamide;
- 1-[4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-yl]-ethanone;
- 2-[4-(4-fluorophenoxy)phenyl]pyrimidine-4-carboxamide;
- 2-[4-(4-fluorophenoxy)phenyl]-4-methylpyrimidine;
- 2-methyl-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid sodium salt;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid methylamide;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid dimethylamide;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid *tert*-butylamide;
- 2-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-4-carboxamide;
- 2-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-4-carboxylic acid;
- 2-(4-phenoxyphenyl)-6-(dimethylamino)pyrimidine-4-carboxylic acid dimethylamide;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid 2-hydroxyethylamide;
- 4-[4-(4-fluorophenoxy)phenyl]pyrimidine-2-carboxylic acid hydroxymethyleneamide;
- 2-(2-hydroxyprop-2-yl)-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
- 4-[4-(2,4-difluorophenoxy)phenyl]pyrimidine-2-carboxylic acid 2-morpholin-4-yl-ethyl amide;
- 2-(4,5-dihydro-1H-imidazol-2-yl)-4-[4-(4-fluorophenoxy)phenyl]-pyrimidine;

D4)

B7

2-(3-pyrazolyl)-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
2-(5-isoxazolyl)-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
2-(1-methyl-3-pyrazolyl)-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
2-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-4-carboxylic acid
methanamide;
3-dimethylamino-1-{4-[4-(4-fluorophenoxy)phenyl]pyrimidin-2-yl}propenone;
2-thiomethyl-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
2-methanesulfonyl-4-[4-(4-fluorophenoxy)phenyl]pyrimidine;
2-[4-(4-chloro-2-fluorophenoxy)phenyl]-4-methyl-pyrimidine;
4-[4-(4-fluorophenoxy)-3-fluorophenyl]pyrimidine-2-carboxamide; or
2-[4-(4-fluorophenoxy)-3-fluorophenyl]pyrimidine-4-carboxamide;
or a pharmaceutically acceptable salt, prodrug or solvate thereof.


Please substitute the following claim 42 for the pending claim 42:

B8)

42. (Once Amended) The compound of claim 41, wherein A₃ is N and A₁ and A₂ are CR₂.


Please substitute the following claim 50 for the pending claim 50:

50. (Once Amended) A pharmaceutical composition, comprising the compound of formula:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:

B9
Sub
C3



Y is or R₇, provided that when Y is R₇, R₁ is aminocarbonyl;

A₁ is N and A₂ and A₃ are CR₂; or A₃ is N and A₁ and A₂ are CR₂;

R₁ is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, C(O)R₈, SO₂R₈, OC(O)NH₂, 2-imidazolyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R₂ is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, amino, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₇ is an optionally substituted alkyl;

R₈ is selected from the group consisting of alkyl, alkenyl, alkynyl, OR₉, amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R₈ is not OR₉ when R₁ is SO₂R₈; wherein

R₉ is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and

X is one of O, S, NH, or CH₂ when Y is other than R₇; or

X is one of O, S, NH, CH₂ or absent when Y is R₇; and a pharmaceutically acceptable carrier or diluent.

Sub
C3

B9


Please add the following claims 59-68:

59. (New) A compound of claim 2, wherein said compound is 2-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-4-carboxamide or a pharmaceutically acceptable salt, prodrug or solvate thereof.


60. (New) A compound of claim 59, which is 2-[4-(4-chloro-2-fluorophenoxy)phenyl]pyrimidine-4-carboxamide.

61. (New) A pharmaceutical composition, comprising the compound of claim 59 or claim 60 and a pharmaceutically acceptable carrier or diluent.

62. (New) A compound of claim 1 having the Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein:



provided that when Y is R₇, R₁ is aminocarbonyl;

A₁ is N and A₂ and A₃ are CR₂; or A₃ is N and A₁ and A₂ are CR₂;

R₁ is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, C(O)R₈, SO₂R₈, OC(O)NH₂, 2-imidazolynyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R₂ is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, amino, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and

D4
B10

aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₇ is an optionally substituted alkyl;

R₈ is selected from the group consisting of alkyl, alkenyl, alkynyl, OR₉, amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R₈ is not OR₉ when R₁ is SO₂R₈; wherein

R₉ is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and


X is one of O, S, NH, or CH₂ when Y is other than R₇; or

X is one of O, S, NH, CH₂ or absent when Y is R₇;


with the proviso that R₂ is not methoxy if R₅ is trifluoromethyl, R₆ is H, X is O and R₁ is SO₂CH₂Ph.

63. (New)

A compound of claim 1 having the Formula I:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:



Y is

or R₇,

provided that when Y is R₇, R₁ is aminocarbonyl;

A₁ is N and A₂ and A₃ are CR₂; or A₃ is N and A₁ and A₂ are CR₂;

R_1 is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, $C(O)R_8$, SO_2R_8 , $OC(O)NH_2$, 2-imidazolyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R_2 is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R_1 and R_2 are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

R_3 , R_4 , R_5 , and R_6 are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R_7 is an optionally substituted alkyl;

R_8 is selected from the group consisting of alkyl, alkenyl, alkynyl, OR_9 , amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R_8 is not OR_9 when R_1 is SO_2R_8 ; wherein

R_9 is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and

X is one of O, S, NH, or CH_2 when Y is other than R_7 ; or

X is one of O, S, NH, CH_2 or absent when Y is R_7 ;

with the proviso that R_2 is not methoxy if R_5 is trifluoromethyl, R_6 is H, X is O and R_1 is SO_2CH_2Ph .

64. (New) The compound of claim 2, wherein A_1 is N and A_2 and A_3 are


CR_2 .

65. (New) The compound of claim 17, wherein A_1 is N and A_2 and A_3 are


CR_2 .

66. (New) The compound of claim 41, wherein A₁ is N and A₂ and A₃ are CR₂.

67. (New) A pharmaceutical composition of claim 50, comprising the compound of formula:



or a pharmaceutically acceptable salt or solvate thereof, wherein:



Y is

or R₇, provided that when Y is R₇, R₁ is

aminocarbonyl;

A₁ is N and A₂ and A₃ are CR₂; or A₃ is N and A₁ and A₂ are CR₂;

R₁ is selected from the group consisting of an optionally substituted alkyl, amino, alkylthio, C(O)R₈, SO₂R₈, OC(O)NH₂, 2-imidazolyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R₂ is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, amino, alkylamino, dialkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₇ is an optionally substituted alkyl;

D4
1310

D41

R_8 is selected from the group consisting of alkyl, alkenyl, alkynyl, OR_9 , amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R_8 is not OR_9 when R_1 is SO_2R_8 ; wherein


R_9 is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and

X is one of O, S, NH, or CH_2 when Y is other than R_7 ; or


1310

X is one of O, S, NH, CH_2 or absent when Y is R_7 ; and a pharmaceutically acceptable carrier or diluent.

68. (New) A pharmaceutical composition of claim 50, comprising the compound of formula:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:



Y is

or R_7 , provided that when Y is R_7 , R_1 is

aminocarbonyl;

A_1 is N and A_2 and A_3 are CR_2 ; or A_3 is N and A_1 and A_2 are CR_2 ;

R_1 is selected from the group consisting an optionally substituted alkyl, amino, alkylthio, $C(O)R_8$, SO_2R_8 , $OC(O)NH_2$, 2-imidazolynyl, 2-imidazolyl, 3-pyrazolyl, 5-isoxazolyl, and 3-(1,2,4)-triazolyl;

each R_2 is selected from the group consisting of hydrogen, optionally substituted alkyl, alkenyl, or alkynyl, halogen, hydroxy, cycloalkyl, cyano, alkylamino,

Sub
C6
diaalkylamino, alkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, and aralkylcarbonylamino; or R₁ and R₂ are taken together with the carbon atoms to which they are attached to form a heterocyclic ring;

Bio
R₃, R₄, R₅, and R₆ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, halogen, haloalkyl, hydroxyalkyl, hydroxy, nitro, amino, cyano, amide, carboxyalkyl, alkoxyalkyl, ureido, acylamino, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₇ is an optionally substituted alkyl;

R₈ is selected from the group consisting of alkyl, alkenyl, alkynyl, OR₉, amino, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, dialkylaminoalkylamino, dialkylaminoalkenylamino, alkylaminoalkenyl-amino, hydroxyaminoalkenylamino, cycloalkyl, heterocycloalkyl, cycloalkylalkylamino, heterocycloalkylamino, aryl, arylalkyl, arylalkenyl, arylalkynyl, and arylalkylamino, all of which can be optionally substituted, provided that R₈ is not OR₉ when R₁ is SO₂R₈; wherein

R₉ is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkali metal; and

X is one of O, S, NH, or CH₂ when Y is other than R₇; or

X is one of O, S, NH, CH₂ or absent when Y is R₇; and a pharmaceutically acceptable carrier or diluent.